ELSEVIER

Neuroscience Letters 307 (2001) 200–203

Neuroscience Letters

www.elsevier.com/locate/neulet

Author Index to Volume 307

Abicht, A., see Jahn, K., 89
Abraham, K.E., Brewer, K.L., Expression of *c-fos* mRNA is increased and related to dynorphin mRNA expression following excitotoxic spinal cord injury in the rat, 187
Abraham, W.C., see Mellentin, C., 13
Adelsberger, H., see Hapfelmeier, G., 97
Adler, T., see Dettmers, C., 109
Ando, H., Sasaki, Y., Okada, H., Urano, A., Prepubertal increases in the levels of two salmon gonadotropin-releasing hormone mRNAs in the ventral telencephalon and preoptic area of masu salmon, 93
Ashur-Fabian, O., Giladi, E., Furman, S., Steingart, R.A., Wollman, Y., Fridkin, M., Brenneman, D.E., Gozes, I., Vasoactive intestinal peptide and related molecules induce nitrite accumulation in the extracellular milieu of rat cerebral cortical cultures, 167
Axer, H., see Krings, T., 192

Bandopadhyay, R., de Silva, R., Khan, N., Graham, E., Vaughan, J., Engelender, S., Ross, C., Morris, H., Morris, C., Wood, N.W., Daniel, S., Lees, A., No pathogenic mutations in the synphilin-1 gene in Parkinson's disease, 125
Bartels, M., see Plewnia, C., 41
Beyer, C., see Ivanova, T., 21
Brückner, L., see Dettmers, C., 109
Brenneman, D.E., see Ashur-Fabian, O., 167
Brewer, K.L., see Abraham, K.E., 187
Bufler, J., see Jahn, K., 89

Carta, G., see Nava, F., 57
Cataudella, T., see Stanzani, S., 135
Chang, C.-K., Lin, M.-T., DL-Tetrahydropalmatine may act through inhibition of amygdaloid release of dopamine to inhibit an epileptic attack in rats, 163
Chen, F.-P., see Hsieh, J.-C., 105
Chen, M.-C., see Hsieh, J.-C., 105
Cheng, H.-C., see Hsieh, J.-C., 105
Cheng, J.-T., see Liu, I.-M., 81
Chennaoui, M., Drogou, C., Gomez-Merino, D., Grimaldi, B., Fillion, G., Guezennec, C.Y., Endurance training effects on 5-HT_{1B} receptors mRNA expression in cerebellum, striatum, frontal cortex and hippocampus of rats, 33
Chi, T.-C., see Liu, I.-M., 81
Chiken, S., Hatanaka, N., Tokuno, H., Distribution of median, ulnar and radial motoneurons in the monkey spinal cord: a retrograde triple-labeling study, 143
Cho, B.-K., see Chung, S.K., 69
Chung, J.-H., see Kim, E.-H., 113
Chung, S.K., Sim, K.B., Cho, B.-K., Hong, S.-K., Wang, K.-C., Is the reclosure of surgically induced open neural tube defect the repetition of primary neurulation? An experimental study with chick embryos, 69
Coenen, V.A., see Krings, T., 192
Cohen, L., see Plewnia, C., 41

Daniel, S., see Bandopadhyay, R., 125
Darlington, C.L., Ross, A., King, J., Smith, P.F., Menstrual cycle effects on postural stability but not optokinetic function, 147
De Nucci, G., see Rogério, F., 61
de Silva, R., see Bandopadhyay, R., 125
de Souza Queiroz, L., see Rogério, F., 61
Deguchi, S., see Mima, K., 117
Dettmers, C., Adler, T., Rzanny, R., van Schayck, R., Gaser, C., Weiss, T., Miltner, W.H., Brückner, L., Weiller, C., Increased excitability in the primary motor cortex and supplementary motor area in patients with phantom limb pain after upper limb amputation, 109
Drogou, C., see Chennaoui, M., 33

Eder, M., see Hapfelmeier, G., 97
Engelender, S., see Bandopadhyay, R., 125
Engelhardt, B., see Wolburg, H., 77
Erwin, J.M., see Hof, P.R., 137

Fillion, G., see Chennaoui, M., 33
Fridkin, M., see Ashur-Fabian, O., 167
Fuchs, E., see Kole, M.H.P., 53
Fujihira, K., see Tabuchi, K., 29
Fukushima, N., see Kato, T., 17
Furman, S., see Ashur-Fabian, O., 167

Gaser, C., see Dettmers, C., 109
Gera, L., see Pan, W., 25
Gerloff, C., see Plewnia, C., 41
Geurts, M., see Robinet, E.A., 45
Giladi, E., see Ashur-Fabian, O., 167
Gilsbach, J.M., see Krings, T., 192
Gomez-Merino, D., see Chennaoui, M., 33
Gozal, D., see Schurr, A., 151
Gozal, E., see Schurr, A., 151
Gozes, I., see Ashur-Fabian, O., 167
Graf von Keyserlingk, D., see Krings, T., 192
Graham, E., see Bandopadhyay, R., 125
Grimaldi, B., see Chennaoui, M., 33
Guezennec, C.Y., see Chennaoui, M., 33

Haase, C.G., Schmidt, S., Detection of brain-specific autoantibodies to myelin oligodendrocyte glycoprotein, S100β and myelin basic protein in patients with Devic's neuromyelitis optica, 131
Hapfelmeier, G., Haseneder, R., Eder, M., Adelsberger, H., Kochs, E., Rammes, G., Zieglgänsberger, W., Isoflurane slows inactivation kinetics

PII: S0304-3940(01)02016-X

ELSEVIER

Neuroscience Letters 307 (2001) 200–203

Neuroscience Letters

www.elsevier.com/locate/neulet

Author Index to Volume 307

Abicht, A., see Jahn, K., 89
Abraham, K.E., Brewer, K.L., Expression of *c-fos* mRNA is increased and related to dynorphin mRNA expression following excitotoxic spinal cord injury in the rat, 187
Abraham, W.C., see Mellentin, C., 13
Adelsberger, H., see Hapfelmeier, G., 97
Adler, T., see Dettmers, C., 109
Ando, H., Sasaki, Y., Okada, H., Urano, A., Prepubertal increases in the levels of two salmon gonadotropin-releasing hormone mRNAs in the ventral telencephalon and preoptic area of masu salmon, 93
Ashur-Fabian, O., Giladi, E., Furman, S., Steingart, R.A., Wollman, Y., Fridkin, M., Brenneman, D.E., Gozes, I., Vasoactive intestinal peptide and related molecules induce nitrite accumulation in the extracellular milieu of rat cerebral cortical cultures, 167
Axer, H., see Krings, T., 192

Bandopadhyay, R., de Silva, R., Khan, N., Graham, E., Vaughan, J., Engelender, S., Ross, C., Morris, H., Morris, C., Wood, N.W., Daniel, S., Lees, A., No pathogenic mutations in the synphilin-1 gene in Parkinson's disease, 125
Bartels, M., see Plewnia, C., 41
Beyer, C., see Ivanova, T., 21
Brückner, L., see Dettmers, C., 109
Brenneman, D.E., see Ashur-Fabian, O., 167
Brewer, K.L., see Abraham, K.E., 187
Bufler, J., see Jahn, K., 89

Carta, G., see Nava, F., 57
Cataudella, T., see Stanzani, S., 135
Chang, C.-K., Lin, M.-T., DL-Tetrahydropalmatine may act through inhibition of amygdaloid release of dopamine to inhibit an epileptic attack in rats, 163
Chen, F.-P., see Hsieh, J.-C., 105
Chen, M.-C., see Hsieh, J.-C., 105
Cheng, H.-C., see Hsieh, J.-C., 105
Cheng, J.-T., see Liu, I.-M., 81
Chennaoui, M., Drogou, C., Gomez-Merino, D., Grimaldi, B., Fillion, G., Guezennec, C.Y., Endurance training effects on 5-HT_{1B} receptors mRNA expression in cerebellum, striatum, frontal cortex and hippocampus of rats, 33
Chi, T.-C., see Liu, I.-M., 81
Chiken, S., Hatanaka, N., Tokuno, H., Distribution of median, ulnar and radial motoneurons in the monkey spinal cord: a retrograde triple-labeling study, 143
Cho, B.-K., see Chung, S.K., 69
Chung, J.-H., see Kim, E.-H., 113
Chung, S.K., Sim, K.B., Cho, B.-K., Hong, S.-K., Wang, K.-C., Is the reclosure of surgically induced open neural tube defect the repetition of primary neurulation? An experimental study with chick embryos, 69
Coenen, V.A., see Krings, T., 192
Cohen, L., see Plewnia, C., 41

Daniel, S., see Bandopadhyay, R., 125
Darlington, C.L., Ross, A., King, J., Smith, P.F., Menstrual cycle effects on postural stability but not optokinetic function, 147
De Nucci, G., see Rogério, F., 61
de Silva, R., see Bandopadhyay, R., 125
de Souza Queiroz, L., see Rogério, F., 61
Deguchi, S., see Mima, K., 117
Dettmers, C., Adler, T., Rzanny, R., van Schayck, R., Gaser, C., Weiss, T., Miltner, W.H., Brückner, L., Weiller, C., Increased excitability in the primary motor cortex and supplementary motor area in patients with phantom limb pain after upper limb amputation, 109
Drogou, C., see Chennaoui, M., 33

Eder, M., see Hapfelmeier, G., 97
Engelender, S., see Bandopadhyay, R., 125
Engelhardt, B., see Wolburg, H., 77
Erwin, J.M., see Hof, P.R., 137

Fillion, G., see Chennaoui, M., 33
Fridkin, M., see Ashur-Fabian, O., 167
Fuchs, E., see Kole, M.H.P., 53
Fujihira, K., see Tabuchi, K., 29
Fukushima, N., see Kato, T., 17
Furman, S., see Ashur-Fabian, O., 167

Gaser, C., see Dettmers, C., 109
Gera, L., see Pan, W., 25
Gerloff, C., see Plewnia, C., 41
Geurts, M., see Robinet, E.A., 45
Giladi, E., see Ashur-Fabian, O., 167
Gilsbach, J.M., see Krings, T., 192
Gomez-Merino, D., see Chennaoui, M., 33
Gozal, D., see Schurr, A., 151
Gozal, E., see Schurr, A., 151
Gozes, I., see Ashur-Fabian, O., 167
Graf von Keyserlingk, D., see Krings, T., 192
Graham, E., see Bandopadhyay, R., 125
Grimaldi, B., see Chennaoui, M., 33
Guezennec, C.Y., see Chennaoui, M., 33

Haase, C.G., Schmidt, S., Detection of brain-specific autoantibodies to myelin oligodendrocyte glycoprotein, S100β and myelin basic protein in patients with Devic's neuromyelitis optica, 131
Hapfelmeier, G., Haseneder, R., Eder, M., Adelsberger, H., Kochs, E., Rammes, G., Zieglgänsberger, W., Isoflurane slows inactivation kinetics

PII: S0304-3940(01)02016-X

of rat recombinant $\alpha_1\beta_2\gamma_{2L}$ $GABA_A$ receptors: enhancement of GABAergic transmission despite an open-channel block, 97
Hara, A., see Tabuchi, K., 29
Hartmann, T., see Schönknecht, P., 122
Haseneder, R., see Hapfelmeier, G., 97
Hatanaka, N., see Chiken, S., 143
Hayashi, Y., see Tanaka, J., 155
Ho, L.-T., see Hsieh, J.-C., 105
Hof, P.R., Nimchinsky, E.A., Perl, D.P., Erwin, J.M., An unusual population of pyramidal neurons in the anterior cingulate cortex of hominids contains the calcium-binding protein calretinin, 139
Höller, M., see Krings, T., 192
Hong, S.-K., see Chung, S.K., 69
Hsieh, J.-C., Tu, C.-H., Chen, F.-P., Chen, M.-C., Yeh, T.-C., Cheng, H.-C., Wu, Y.-T., Liu, R.-S., Ho, L.-T., Activation of the hypothalamus characterizes the acupuncture stimulation at the analgesic point in human: a positron emission tomography study, 105
Huda, K., Salunga, T.L., Matsunami, K., Dopaminergic inhibition of excitatory inputs onto pyramidal tract neurons in cat motor cortex, 175

Ikegaya, Y., see Nakao, K., 159
Ishida, Y., Shirokawa, T., Komatsu, Y., Isobe, K., Changes in cortical noradrenergic axon terminals of locus coeruleus neurons in aged F344 rats, 197
Ishihara, K., Sasa, M., Potentiation of $5\text{-}HT_3$ receptor functions in the hippocampal CA1 region of rats following repeated electroconvulsive shock treatments, 37
Isobe, K., see Ishida, Y., 197
Ivanova, T., Beyer, C., Pre- and postnatal expression of brain-derived neurotrophic factor mRNA/protein and tyrosine protein kinase receptor B mRNA in the mouse hippocampus, 21
Izumi, Y., see Nishimura, M., 128

Jahn, K., Mohammadi, B., Krampfl, K., Abicht, A., Lochmüller, H., Bufler, J., Deactivation and desensitization of mouse embryonic- and adult-type nicotinic receptor channel currents, 89
Jang, M.-H., see Kim, E.-H., 113
Jensen, M., see Schönknecht, P., 122

Kaji, R., see Nishimura, M., 128
Kastin, A.J., see Pan, W., 25
Kato, T., Yokouchi, K., Fukushima, N., Kawagishi, K., Li, Z., Moriizumi, T., Continual replacement of newly-generated olfactory neurons in adult rats, 17
Kawagishi, K., see Kato, T., 17
Kawakami, H., see Nishimura, M., 128
Khan, N., see Bandopadhyay, R., 125
Kim, C.-J., see Kim, E.-H., 113
Kim, E.-H., Kim, Y., Jang, M.-H., Lim, B.-V., Kim, Y.-J., Chung, J.-H., Kim, C.-J., Auricular acupuncture decreases neuropeptide Y expression in the hypothalamus of food-deprived Sprague–Dawley rats, 113
Kim, Y., see Kim, E.-H., 113
Kim, Y.-J., see Kim, E.-H., 113
Kimura, K., see Okamoto, S., 179
King, J., see Darlington, C.L., 147
Klein, T., see Town, T., 101
Klimaschewski, L., Increased innervation of rat preganglionic sympathetic neurons by substance P containing nerve fibers in response to spinal cord injury, 73
Klinga, K., see Schönknecht, P., 122
Kobayashi, M., see Narita, M., 85
Kochs, E., see Hapfelmeier, G., 97
Kofler, M., Müller, J., Reggiani, L., Valls-Solé, J., Influence of age on auditory startle responses in humans, 65
Kohsaka, S., see Nakajima, K., 171
Kole, M.H.P., Koolhaas, J.M., Luiten, P.G.M., Fuchs, E., High-voltage-activated Ca^{2+} currents and the excitability of pyramidal neurons in the hippocampal CA3 subfield in rats depend on corticosterone and time of day, 53
Komatsu, Y., see Ishida, Y., 197
Koolhaas, J.M., see Kole, M.H.P., 53
Krampfl, K., see Jahn, K., 89
Krausz, G., see Pfurtscheller, G., 49
Krings, T., Coenen, V.A., Axer, H., Reinges, M.H.T., Höller, M., Graf von Keyserlingk, D., Gilsbach, J.M., Thron, A., In vivo 3D visualization of normal pyramidal tracts in human subjects using diffusion weighted magnetic resonance imaging and a neuronavigation system, 192
Kudo, N., see Nakayama, K., 1
Kuno, S., see Nishimura, M., 128
Kurihara, T., see Nakajima, K., 171
Kusakari, J., see Tabuchi, K., 29

Langone, F., see Rogério, F., 61
Lees, A., see Bandopadhyay, R., 125
Li, Y.-Q., see Wu, S.-X., 183
Li, Z., see Kato, T., 17
Liebner, S., see Wolburg, H., 77
Lim, B.-V., see Kim, E.-H., 113
Lin, M.-T., see Chang, C.-K., 163
Lin, M.-T., see Liu, I.-M., 81
Liu, I.-M., Chi, T.-C., Shiao, G.-C., Lin, M.-T., Cheng, J.-T., Loss of plasma glucose lowering response to cold stress in opioid μ-receptor knock-out diabetic mice, 81
Liu, R.-S., see Hsieh, J.-C., 105
Lochmüller, H., see Jahn, K., 89
Luiten, P.G.M., see Kole, M.H.P., 53

Müller, J., see Kofler, M., 65
Maestre, G.E., see Molero, A.E., 5
Maloteaux, J.-M., see Robinet, E.A., 45
Maruyama, H., see Nishimura, M., 128
Matsuki, N., see Nakao, K., 159
Matsunami, K., see Huda, K., 175
Mellentin, C., Abraham, W.C., Priming stimulation of group II metabotropic glutamate receptors inhibits the subsequent induction of rat hippocampal long-term depression in vitro, 13
Miltner, W.H., see Dettmers, C., 109
Mima, K., Deguchi, S., Yamauchi, T., Characterization of 5′ flanking region of α isoform of rat Ca^{2+}/calmodulin-dependent protein kinase II gene and neuronal cell type specific promoter activity, 117
Miyakubo, H., see Tanaka, J., 155
Mohammadi, B., see Jahn, K., 89
Molero, A.E., Pino-Ramírez, G., Maestre, G.E., Modulation by age and gender of risk for Alzheimer's disease and vascular dementia associated with the apolipoprotein E-ε4 allele in Latin Americans: findings from the Maracaibo Aging Study, 5
Moriizumi, T., see Kato, T., 17
Morris, C., see Bandopadhyay, R., 125
Morris, H., see Bandopadhyay, R., 125
Mullan, M., see Town, T., 101

Nakajima, K., Tohyama, Y., Kohsaka, S., Kurihara, T., Ability of rat microglia to uptake extracellular glutamate, 171
Nakamura, S., see Nishimura, M., 128
Nakao, K., Ikegaya, Y., Yamada, M.K., Nishiyama, N., Matsuki, N., Spatial performance correlates with long-term potentiation of the dentate gyrus but not of the CA1 region in rats with fimbria–fornix lesions, 159
Nakayama, K., Nishimaru, H., Kudo, N., Developmental changes in 5-hydroxytryptamine-induced rhythmic activity in the spinal cord of rat fetuses in vitro, 1
Narita, M., see Narita, M., 85
Narita, M., Tamaki, H., Kobayashi, M., Soma, M., Narita, M., Suzuki, T., Changes in Ca^{2+}-dependent protein kinase C isoforms induced by chronic ethanol treatment in mice, 85
Nava, F., Carta, G., Melatonin reduces anxiety induced by lipopolysaccharide in the rat, 57
Neuper, C., see Pfurtscheller, G., 49
Nimchinsky, E.A., see Hof, P.R., 139
Nishimaru, H., see Nakayama, K., 1
Nishimura, M., Kawakami, H., Maruyama, H., Izumi, Y., Kuno, S., Kaji, R., Nakamura, S., Influence of interleukin-1β gene polymorphism on age-at-onset of spinocerebellar ataxia 6 (SCA6) in Japanese patients, 128
Nishiyama, N., see Nakao, K., 159

Obregon, D., see Town, T., 101
Okada, H., see Ando, H., 93
Okamoto, S., Kimura, K., Saito, M., Anorectic effect of leptin is mediated by hypothalamic corticotropin-releasing hormone, but not by urocortin, in rats, 179
Okubo, H., see Tabuchi, K., 29
Okumura, T., see Tanaka, J., 155

Pan, W., Kastin, A.J., Gera, L., Stewart, J.M., Bradykinin antagonist decreases early disruption of the blood–spinal cord barrier after spinal cord injury in mice, 25
Pantel, J., see Schönknecht, P., 122
Pauwels, P.J., see Robinet, E.A., 45
Payne, R.S., see Schurr, A., 151
Pellitteri, R., see Stanzani, S., 135
Perl, D.P., see Hof, P.R., 139
Pfurtscheller, G., Woertz, M., Krausz, G., Neuper, C., Distinction of different fingers by the frequency of stimulus induced beta oscillations in the human EEG, 49
Pino-Ramírez, G., see Molero, A.E., 5
Plewnia, C., Bartels, M., Cohen, L., Gerloff, C., Noradrenergic modulation of human cortex excitability by the presynaptic α_2-antagonist yohimbine, 41

Rammes, G., see Hapfelmeier, G., 97
Reggiani, L., see Kofler, M., 65
Reinges, M.H.T., see Krings, T., 192
Robinet, E.A., Geurts, M., Maloteaux, J.-M., Pauwels, P.J., Chronic treatment with certain antipsychotic drugs preserves upregulation of regulator of G-protein signalling 2 mRNA in rat striatum as opposed to c-fos mRNA, 45
Rogério, F., Teixeira, S.A., de Souza Queiroz, L., De Nucci, G., Langone, F., Expression of neuronal isoform of nitric oxide synthase in spinal neurons of neonatal rats after sciatic nerve transection, 61
Ross, A., see Darlington, C.L., 147
Ross, C., see Bandopadhyay, R., 125
Russo, A., see Stanzani, S., 135
Rzanny, R., see Dettmers, C., 109

Saito, M., see Okamoto, S., 179
Sakamaki, K., see Tanaka, J., 155
Salbach, B., see Schönknecht, P., 122
Salunga, T.L., see Huda, K., 175
Sansone, N., see Town, T., 101
Sasa, M., see Ishihara, K., 37
Sasaki, Y., see Ando, H., 93
Schmidt, S., see Haase, C.G., 131
Schönknecht, P., Pantel, J., Klinga, K., Jensen, M., Hartmann, T., Salbach, B., Schröder, J., Reduced cerebrospinal fluid estradiol levels are associated with increased β-amyloid levels in female patients with Alzheimer's disease, 122
Schröder, J., see Schönknecht, P., 122
Schurr, A., Payne, R.S., Tseng, M.T., Gozal, E., Gozal, D., Excitotoxic preconditioning elicited by both glutamate and hypoxia and abolished by lactate transport inhibition in rat hippocampal slices, 151
Shiao, G.-C., see Liu, I.-M., 81
Shirokawa, T., see Ishida, Y., 197
Sim, K.B., see Chung, S.K., 69
Smith, P.F., see Darlington, C.L., 147
Soma, M., see Narita, M., 85
Stanzani, S., Russo, A., Pellitteri, R., Storaci, G., Cataudella, T., Branching projections of catecholaminergic ventrolateral reticular neurons to the fastigial nucleus and superior colliculus in the rat: triple labelling procedure, 135
Steingart, R.A., see Ashur-Fabian, O., 167
Stewart, J.M., see Pan, W., 25
Storaci, G., see Stanzani, S., 135
Suzuki, T., see Narita, M., 85

Tabira, T., see Tanahashi, H., 9
Tabuchi, K., Okubo, H., Fujihira, K., Tsuji, S., Hara, A., Kusakari, J., Protection of outer hair cells from reperfusion injury by an iron chelator and a nitric oxide synthase inhibitor in the guinea pig cochlea, 29
Tamaki, H., see Narita, M., 85
Tan, J., see Town, T., 101
Tanahashi, H., Tabira, T., Three novel alternatively spliced isoforms of the human beta-site amyloid precursor protein cleaving enzyme (BACE) and their effect on amyloid beta-peptide production, 9
Tanaka, J., Miyakubo, H., Okumura, T., Sakamaki, K., Hayashi, Y., Estrogen decreases the responsiveness of subfornical organ neurons projecting to the hypothalamic paraventricular nucleus to angiotensin II in female rats, 155
Teixeira, S.A., see Rogério, F., 61
Thron, A., see Krings, T., 192
Tohyama, Y., see Nakajima, K., 171
Tokuno, H., see Chiken, S., 143
Town, T., Tan, J., Sansone, N., Obregon, D., Klein, T., Mullan, M., Characterization of murine immunoglobulin G antibodies against human amyloid-β_{1-42}, 101
Tseng, M.T., see Schurr, A., 151
Tsuji, S., see Tabuchi, K., 29
Tu, C.-H., see Hsieh, J.-C., 105

Urano, A., see Ando, H., 93

Valls-Solé, J., see Kofler, M., 65
van Schayck, R., see Dettmers, C., 109

Vaughan, J., see Bandopadhyay, R., 125

Wang, K.-C., see Chung, S.K., 69
Wang, W., see Wu, S.-X., 183
Wang, Y.-Y., see Wu, S.-X., 183
Weiller, C., see Dettmers, C., 109
Weiss, T., see Dettmers, C., 109
Woertz, M., see Pfurtscheller, G., 49
Wolburg, H., Wolburg-Buchholz, K., Liebner, S., Engelhardt, B., Claudin-1, claudin-2 and claudin-11 are present in tight junctions of choroid plexus epithelium of the mouse, 77
Wolburg-Buchholz, K., see Wolburg, H., 77
Wollman, Y., see Ashur-Fabian, O., 167
Wood, N.W., see Bandopadhyay, R., 125
Wu, S.-X., Zhu, M., Wang, W., Wang, Y.-Y., Li, Y.-Q., Yew, D.T., Changes of the expression of 5-HT receptor subtype mRNAs in rat dorsal root ganglion by complete Freund's adjuvant-induced inflammation, 183
Wu, Y.-T., see Hsieh, J.-C., 105

Yamada, M.K., see Nakao, K., 159
Yamauchi, T., see Mima, K., 117
Yeh, T.-C., see Hsieh, J.-C., 105
Yew, D.T., see Wu, S.-X., 183
Yokouchi, K., see Kato, T., 17

Zhu, M., see Wu, S.-X., 183
Zieglgänsberger, W., see Hapfelmeier, G., 97